
## FRUIT DOTS Peter Culley



## Fruit Dots




## Canadian Cataloguing in Publication Data

Culley, Peter, 1958-Fruit dots

> Poems. ISBN 0-921331-01-0

I. Title.
PS8555.U44F7 1986 C811'.54 C86-091585-9
PR9199.3.C84F7 1986



in May. As it unfolds
spores are discharged.
Tuft of rusty wool

above the fruiting pinnae,
shallow vase formed, flood
the woods with golden light.

Other osmundas, absence of the fertile fronds. Erect, curving outward,

oblong-lanceolate; obtuse divisions
taller than the sterile
entire or toothed.

Of the midrib, indisium delicate-little fronds, ecstatic presence
of the deeper woods.

Arched across the rock-broken stream, carpeted with low; where precious red-cupped mosses

stretch basswood overhead cliff which forms.

Early meadow rue --

lobed and rounded leaflet

crowds to the roadside

moist hollows: fronds

tremulous on their black

chance to be driving

by a bank overgrown,

posessed by a tormenting love sadly state but firm; aloofness which adds

haunts are dim.

Summit of the slender black and polished, recurved

branches bearing
several slender,
under margin of a lobe

dotted with fragrant heaps.

Entered the cool shade, beauty and luxuriance,

rich green arched above rigid fruit-clusters.

Crown firmed rigid;

by means of underground runners

crawled under

a tight board fence --

free veins with simple veinlets

do not reach

the midvein. Reflexed marginal

rows of bright brown sporangia

wedged in neighboring

crannies, sprang from limestone cliffs;

so blended, with their rocky self-assurance at variance.

Prostrate among the rocks

rise the firm

graceful crowns:

conspicuously marginal

fruit-dots, caudex covered by the bases-apex and the tips
of its pinnules, plumes

of departing summer, of the greenhouse.

Above the black leaf-mould,

immortal in the swamp herbage,

centre shades away

less chaffy stalks; mossy ledges


above star-like stumps of decaying.

A projecting shelf of bluish-green;

leathery, tapering,

decayed at its tip

a flower yet unnamed.



Day count.

Columbine flings out

just uncurled

roots of some great

nodding blossom,

among which the stream forces,

flavored with the essence

once more.

Fastened underneath

into jagged, spreading --

were it not for the difference twice to thrice-pinnate

finally appear naked,

early withering and exposing

beneath a reflexed portion.

Green takes on

a dark heraldic eagle
in a cross-section of the young stalk:

does not wither away,

grown by reason of the licking.

Shall shellfish

in the Andes at fourteen feet
widely spreading at the summit?
Reflexed edge, somewhat open

borne in a continuous line;
stout stalk, from the trunks
somewhat lustrous --


upward turning
on their upper
through the winter.

In rocky woods

having shiny scales,

convex; persistent

along the roadside and in wet, found of the tiniest



its creeping rootstock rise.

At times wear very light, standing brown and dry

from early frosts --

form midway

in masses -- in wet meadows

inappropriateness, wavy-toothed divisions; berry-like bodies usually heart-shaped.

Scattered on the lower surface shale, and conglomerate near limestone cliffs

sprays of the bulblet,
turreted summits.

Rigid grace, slim in crowded ranks,

into moist ravines

necessitates the twisting of its stalk. Shelves of shaded rocks:

surprise--combination of delicacy,

black stemmed

under groups of red, with blackish

and shining, silvery till maturity,

base auricled;

the tangle of wild

close to the roadside fence.

Of some amber-coloured brook the wild bergamot

its swift noiseless flow.

Uncurl in May -present a rather blotched

delicacy, scarlet cluster,

background. Of grain shine
between umbrella-like Brakes:

so recently unfolded texture seems a trifle.

Close against the rails;

as the curves and dimples,
little undulations
of the August woods' border--

